Induction of Labor Checklist



For Obstetrical and Medically Necessary Induction of Labor:

•	onal age (The need to de everity of condition)	liver at a gestational age less than 39 weeks is
•	the following indications	s
□41+0 v □Abrup □Preecl □Gestat □GDM □PROM □Fetal I □Coagu □Pulmo □Choric □Unstat □Other □IUGR □Isoimr □Fetal r □Multip □Twins	weeks tio placentae ampsia ional HTN I Demise lopathy/Thrombophilia mary disease camnionitis ble Lie Fetal compromise munization malformation bles w/ complications w/o complication on, confirm necessity for	Heart disease Liver disease (e.g. cholestasis of pregnancy.) Chronic HTN Diabetes (Type I or II) Renal disease Oligohydramnios Other
(c		consultant name)
Confirm graBe aware of□Pa□Ri□Di		action is planned

- Confirm favorable cervix by Bishops score (See table)
 - ☐Bishop's score >/= 8 for nullipara
 - ☐ Bishop's score >/= 6 for multipara

Bishop's Score Calculation						
Parameter	0	1	2	3		
Dilation (cm)	0	1 - 2	3 - 4	5 - 6		
Effacement, %	0 - 30	40 - 50	60 - 70	≥80		
Station (-3 to +3)	- 3	-2	-1, 0	≥+1		
Consistency	Firm	Medium	Soft			
Position	Posterior	Middle	Anterior			
ACOG Patient Safety Checklist No. 5; December, 2011						

For all Inductions:

- Provide patient with written educational material on induction of labor
- Obtain signed induction of labor education form
- Remind patient to call Labor and Delivery (or designee) prior to leaving home on the day of the induction

References:

ACOG Committee Opinion, No.560, 2013

ACOG Patient Safety Checklist No 2. Inpatient Induction of Labor December 2011, reaffirmed 2014