#### 2. Hospital & QI Team Overview


Yorleni Ambrosio, CNS/APN, Neo Team Lead Aaron Muller, MD, Physician Team Lead Michelle Wrobel, RN, Nurse Lead Rachel Shapiro, NICU Social Worker Guy Steinberg, MD, Maternal Fetal Medicine



#### 3. MNO-Neo Data


Breastfeeding


100% of OEN cases in 2019 received maternal breastmilk at discharge!



Pharm/Non-Pharm Treatments 80% required pharmacologic treatment from 2018-present (area for improvement)

**Coordinated Discharge Plans** 83% of babies met the Goal!






# Advocate Good Samaritan Hospital Advocate Good Samaritan Hospital


### 4. MNO-Neo Progress

**NAS Assessment Tools** ESC method is our primary source of assessment. Mothers are engaged in daily rounds where the ESC assessments are reviewed and are encouraged to utilize the bedside care diary


#### NAS: A Parent's Guide Folder



#### **MNO Team Process**



Screening and Planning



- Prenatal Consult with
- Neonatologist
- "NAS: A Parent's

Baby's Name:			Baby's MRN:					Date:		
Time of baby's feed (start to finish)	Breastfeeding (minutes)	Bottle feeding (total mL)	Time baby fell asleep leading up to this feed	Time baby woke up leading up to this feed	Did the baby feed for at least 10 minutes at breast OR take in at least 10 mL by bottle?	Did the baby sleep for at least 1 hour?	Did the baby console within 10 minutes of being upset? Describe.	Check for pee	Check for poop	Other comments / non- pharmacological care provided
Example 1:30 pm- 1:55 pm	Example L: 10 mins R: 12 mins	Example NA	Example 11:15am	Example 1:15pm	Example Yes	Example Yes	Example Yes. Baby woke up and was difficult to console for ~5 minutes. Changed dirty diaper and then baby latched	Example X	Example X	Example Performed skin to skin following feeding
Example 4:40pm- 4:55pm	Example L: 5 mins R: 0 mins	Example NA	Example 1:15pm	Example 4:30pm	Example No—baby had difficulty latching. Notified RN.	Example Yes	Example No. Attempted skin to skin contact and swaddling. Baby was consoled within 15 mins.	Example X	Example	Example Notified RN about difficulty latching and difficulty with consoling. Want to see lactation consultant.

**Education Campaign** 

## IPGC 2020Neo Storyboard 5. Coordinated Discharge

#### **Discharge and Follow-Up Team Approach**

- home



#### **Plan to Overcome Barriers**

Continue to utilize the ESC assessment tool to reduce the need for pharmacologic tx

Discontinuing use of Finnegan scoring tool as a supplemental resource

Linkage to resources for mom/baby, facilitation of safe home environment Coordination with infant's PCP and ensuring a safe discharge plan Developmental f/u, WIC, therapies, and NAS education for caring for infant at

Utilizing a discharge tool

