

OBSTETRICS

Reduction in racial disparities in severe maternal morbidity from hemorrhage in a large-scale quality improvement collaborative

Elliott K. Main, MD; Shen-Chih Chang, MS, PhD; Ravi Dhurjati, MS, PhD; Valerie Cape, BA; Jochen Profit, MD, MPH; Jeffrey B. Gould, MD, MPH

BACKGROUND: Eliminating persistent racial/ethnic disparities in maternal mortality and morbidity is a public health priority. National strategies to improve maternal outcomes are increasingly focused on quality improvement collaboratives. However, the effectiveness of quality collaboratives for reducing racial disparities in maternity care is understudied.

OBJECTIVE: To evaluate the impact of a hemorrhage quality-improvement collaborative on racial disparities in severe maternal morbidity from hemorrhage.

STUDY DESIGN: We conducted a cross-sectional study from 2011 to 2016 among 99 hospitals that participated in a hemorrhage quality improvement collaborative in California. The focus of the quality collaborative was to implement the national maternal hemorrhage safety bundle consisting of 17 evidence-based recommendations for practice and care processes known to improve outcomes. This analysis included 54,311 women from the baseline period (January 2011 through December 2014) and 19,165 women from the postintervention period (October 2015 through December 2016) with a diagnosis of obstetric hemorrhage during delivery hospitalization. We examined whether racial/ethnic-specific severe maternal morbidity rates in these women with obstetric hemorrhage were reduced from the baseline to the postintervention period. In addition, we conducted Poisson Generalized Estimating Equation models to estimate relative risks and 95% confidence intervals for severe maternal morbidity comparing each racial/ethnic group with white.

RESULTS: During the baseline period, the rate of severe maternal morbidity among women with hemorrhage was 22.1% (12,002/54,311) with the greatest rate observed among black women (28.6%, 973/3404), and the lowest among white women (19.8%, 3124/15,775). The

overall rate fell to 18.5% (3553/19,165) in the postintervention period. Both black and white mothers benefited from the intervention, but the benefit among black women exceeded that of white women (9.0% vs 2.1% absolute rate reduction). The baseline risk of severe maternal morbidity was 1.34 times greater among black mothers compared with white mothers (relative risk, 1.34; 95% confidence interval, 1.27–1.42), and it was reduced to 1.22 (1.05–1.40) in the postintervention period. Sociodemographic and clinical factors explained a part of the black–white differences. After controlling for these factors, the black–white relative risk was 1.22 (95% confidence interval, 1.15–1.30) at baseline and narrowed to 1.07 (1.92–1.24) in the postintervention period. Results were similar when excluding severe maternal morbidity cases with transfusion alone. After accounting for maternal risk factors, the black–white relative risk for severe maternal morbidity excluding transfusion alone was reduced from a baseline of 1.33 (95% confidence interval, 1.16–1.52) to 0.99 (0.76–1.29) in the postintervention period. The most important clinical risk factor for disparate black rates for both severe maternal morbidity and severe maternal morbidity excluding transfusion alone was cesarean delivery, potentially providing another opportunity for quality improvement.

CONCLUSION: A large-scale quality improvement collaborative reduced rates of severe maternal morbidity due to hemorrhage in all races and reduced the performance gap between black and white women. Improving access to highly effective treatments has the potential to decrease disparities for care-sensitive acute hospital-focused morbidities.

Key words: hemorrhage, maternal morbidity, maternal safety, perinatal quality collaboratives, quality improvement, racial/ethnic disparities

Persistent racial/ethnic disparities in maternal mortality and morbidity exist in the United States.^{1–3} Black women continue to be 3–4 times more likely than white women to die during

childbirth.² Severe maternal morbidity (SMM) is a composite measure developed by the Centers for Disease Control and Prevention that includes diagnosis and procedure codes reflecting major complications in childbirth, such as pulmonary edema, renal failure, disseminated intravascular coagulation, hysterectomy, and transfusion.⁴ SMM is 50–100 times more common than maternal death, affecting nearly 60,000 women each year in the United States.^{5,6} The risk of SMM for black women is twice that of white women, even after adjusting for sociodemographic factors and comorbidities.⁷ It is

estimated that more than one half of cases of maternal mortality and morbidity are preventable^{8–11} and could be sensitive to quality of care provided at delivery.^{12,13} Hemorrhage is the most common major complication of childbirth,^{14–17} the most preventable cause of maternal mortality,⁸ and by far the most frequent cause of SMM.¹⁸

Quality-improvement (QI) interventions may reduce disparities only if they improve quality of care and simultaneously reduce the performance gap between racial/ethnic groups.¹⁹ Disparity reduction requires that vulnerable groups with a history of

Cite this article as: Main EK, Chang S-C, Dhurjati R, et al. Reduction in racial disparities in severe maternal morbidity from hemorrhage in a large-scale quality improvement collaborative. *Am J Obstet Gynecol* 2020

0002-9378/\$36.00

© 2020 Elsevier Inc. All rights reserved.

<https://doi.org/10.1016/j.ajog.2020.01.026>



Click Supplemental Materials
under article title in Contents at ajog.org

AJOG at a Glance

Why was the study conducted?

It was not known whether a large-scale quality improvement collaborative could reduce racial disparity in severe maternal complications following obstetric hemorrhage.

Key Findings

In this cross-sectional study that included 73,476 women with obstetric hemorrhage from 99 hospitals who participated in a hemorrhage quality improvement collaborative, the rate of severe maternal morbidity was reduced for all races. The black–white differences were no longer significant following case mix adjustment.

What does this add to what is known?

Maternal quality-improvement activities that focus on improving access to highly effective treatments have the potential to reduce racial disparities for care-sensitive acute hospital-focused morbidities such as hemorrhage.

worse outcomes either receive a greater degree of benefit from the quality intervention or have greater access to the intervention. Otherwise, improvement efforts may compound existing disparities by preferentially advantaging white populations.^{20–23} Unfortunately, little is known about the impact of QI interventions on racial disparities in maternal outcomes.²⁴

Our previous work²⁵ demonstrated that a large-scale QI collaborative resulted in a significant reduction (>20%) in hemorrhage-related SMM following the implementation of a national hemorrhage safety bundle,¹⁸ whereas a comparison set of hospitals, not implementing the hemorrhage safety bundle, remained unchanged. In this report, we examine whether the QI collaborative would be able to reduce the gap between black and white rates of SMM from hemorrhage.

Materials and Methods

A multihospital quality collaborative focused on improving outcomes from obstetric hemorrhage was offered to all California hospitals.²⁵ Ninety-nine hospitals averaging 250,000 annual births choose to participate. The collaborative, led by the California Maternal Quality Care Collaborative (CMQCC), began in January 2015 with intensive activities lasting for 18 months. This report includes 6

additional months of collaborative data not presented in an earlier report²⁵ and an analysis of results by race and ethnicity.

The emphasis of the quality collaborative was to implement the national maternal hemorrhage safety bundle consisting of 17 evidence-based recommendations for practice and care processes known to improve outcomes (Table 1).¹⁸ The implementation strategy was an adaptation of the Institute for Health Care Improvement collaborative model creating a community of learning, including 2 participant face-to-face meetings, and monthly check-in calls. Hospitals were organized into small teams of 6–8 hospitals led by physician and nurse mentors who provided QI coaching.²⁶ This involved monthly team support and advice for the assessment of barriers and improvement strategies. Baseline outcome data were collected for 48 months from January 2011 through December 2014. The postintervention period was from October 2015 to December 2016. We compared baseline outcome measures with those collected in the postintervention period to examine the effect of the intervention on SMM among the racial/ethnic groups.

There were 977,968 deliveries in the 4-year baseline period and 314,750 deliveries in the postintervention period, representing one half of all births in California. Obstetric hemorrhage was

identified in approximately 6% of women in both time periods (56,865 at baseline and 20,278 during the post-intervention period). Obstetric hemorrhage was defined as patients with *International Statistical Classification of Diseases, Ninth and Tenth Revision* diagnosis codes for antepartum or postpartum hemorrhage, placenta previa, and abruption placentae. Hemorrhage often is undercoded, which can be largely corrected by the addition of procedure codes for transfusion, given the very low rate of transfusion for other indications (codes are provided in Supplemental Table 1).

Discharge diagnosis and procedure codes were obtained from the CMQCC California Maternal Data Center. CMQCC uses a modified form of a previously published probabilistic algorithm to link maternal and newborn hospital discharge records with birth certificates.²⁷ Linkage rates routinely exceed 98%. The baseline hemorrhage population (56,865) in this study was slightly lower than reported in the initial study²⁵ (57,320) after excluding 455 (0.8%) because of nonlinkage to birth certificates resulting in missing socio-demographic factors. Birth certificates were received from the California Department of Public Health 45 days after the end of each month. Discharge files were received from the Office of Statewide Health Planning and Development on a semiannual basis (delayed by 6–9 months). Institutional review board approval was obtained from Stanford University as the study host and the California Committee for the Protection of Human Subjects for the use of the state data sets.

Self-reported race and ethnicity were obtained from birth certificates. All women were categorized into 1 of 3 ethnic groups (Hispanic, non-Hispanic, and unknown/missing), and 1 of 7 racial groups (white, black, Asian, Pacific Islander, American Indian, other, and unknown). We collapsed the “Pacific Islander” and “American Indian” groups with the “other” category and created a 5-category race/ethnicity measure: Hispanic, non-Hispanic white (white), non-Hispanic black (black),

TABLE 1

California Partnership for Maternal Safety Collaborative: hemorrhage safety bundle elements^a

Readiness domain

Hemorrhage cart/including instructions
cards for intrauterine balloons and
compression stitches

Immediate access to hemorrhage
medications (kit or equivalent)

Hemorrhage response team established
(anesthesia, blood bank, advanced
gynecological surgery, and other services)

Massive transfusion protocol established

Emergency release protocol established
for O-negative and uncross-matched
units of red blood cells

Protocol established for those who refuse blood products

Unit education to hemorrhage protocols

Regular unit-based drills with debriefs for
obstetric hemorrhage

Recognition and prevention domain

Assessment of hemorrhage risk (prenatal,
admission, and other)

Measurement of cumulative blood loss
(formal and as quantitative as possible)

Active management of third stage of labor
(standard protocol for oxytocin at birth)

Response domain

Use of unit-standard, stage-based obstetric
hemorrhage emergency management plan
with checklists

Support program for patients, families, and staff for all major obstetric hemorrhages

Reporting and systems learning domain

Establish culture of huddles to plan for high-risk patients

Post-event debriefing to quickly assess what went well and what could have been improved

Multidisciplinary reviews of all serious
hemorrhages for system issues

Monitor outcomes and progress in process measures in perinatal quality improvement committee

^a Based on the Alliance for Innovation in Maternal Health (AIM): Consensus Bundle on Obstetric Hemorrhage¹⁸ and condensed from a previous report.²⁵

Main et al. Reduction in racial disparities in severe maternal morbidity from hemorrhage in a large-scale quality improvement collaborative. *Am J Obstet Gynecol* 2020.

Asian, and others. In total, 4.8% of women had an unknown or missing race or ethnicity and were removed from the analysis. The final sample for analysis consisted of 54,311 women in the baseline and 19,165 women in the postintervention period.

The main outcome measure was the rate of SMM among women diagnosed with hemorrhage. Corresponding *International Statistical Classification of*

Diseases codes for SMM are listed on the Centers for Disease Control and Prevention website.⁴ Transfusion is the most common morbidity for SMM. To identify the effect on other morbidities, we also evaluated rates of SMM excluding transfusion-only cases. As transfusion is also part of the definition for hemorrhage (widely used in all state collaboratives in the AIM project and in the prior report),²⁵ the addition of

SMM without transfusion in the numerator provides additional perspective.

We considered the following risk factors for SMM from obstetric hemorrhage: mother's sociodemographic characteristics (maternal age, education, parity, and insurance status), clinical factors (number of prenatal visits, pre-pregnancy body mass index, multiple pregnancy, chronic hypertension,

TABLE 2

Maternal characteristics of Hemorrhage Quality Collaborative Participants and California delivery population, baseline period: 2011–2014

	Collaborative participants n (%)	All California n (%)
Patient characteristics		
Number of deliveries	977,968	1,888,173
Race/ethnicity		
White	309,837 (32.7)	539,396 (29.2)
Black	55,548 (5.9)	106,530 (5.8)
Asian	129,833 (13.7)	216,504 (11.7)
Hispanic	406,626 (42.9)	912,675 (49.3)
Other	45,142 (4.8)	74,678 (4.0)
Missing	30,982	38,390
Age at delivery, y		
<18	15,384 (1.6)	37,752 (2.0)
18–25	246,082 (25.2)	547,769 (29.0)
26–35	554,126 (56.7)	1,018,273 (53.9)
≥36	162,376 (16.6)	284,379 (15.1)
Education		
Some high school or less	133,389 (14.4)	356,203 (19.6)
High school/GED	206,235 (22.3)	461,488 (25.5)
Some college	262,083 (28.4)	477,368 (26.3)
College grad or more	322,211 (34.9)	518,204 (28.6)
Missing	54,050	74,910
Insurance		
Medicaid/other government-sponsored	364,071 (37.2)	913,050 (48.4)
Private insurance	584,349 (59.8)	916,782 (48.6)
Self-pay/no insurance	29,526 (3.0)	58,310 (3.1)
Missing	22	31
Nulliparous	410,607 (42.0)	752,589 (39.9)
Prepregnancy BMI		
<18.5	19,265 (2.1)	38,587 (2.2)
18.5–25	410,011 (44.9)	767,776 (42.8)
25–30	254,980 (27.9)	512,402 (28.6)
30–35	130,985 (14.3)	271,290 (15.1)
35–40	58,705 (6.4)	121,518 (6.8)
≥40	39,347 (4.3)	80,951 (4.5)
Missing	64,675	95,649

Main et al. Reduction in racial disparities in severe maternal morbidity from hemorrhage in a large-scale quality improvement collaborative. Am J Obstet Gynecol 2020. (continued)

Statistical analysis

We used the χ^2 test to examine whether the distributions of maternal social demographic and clinical factors are different between race/ethnic groups and whether they are different in maternal cohorts in the baseline and postintervention period. We then assessed the risk of SMM among women with obstetric hemorrhage by study period and by race/ethnicity. Specifically, we constructed Poisson generalized estimating equation models with sandwich error estimation to estimate relative risks (RRs) and 95% confidence intervals (CIs) for SMM. A estimating equation is a population-average model that accounts for within-hospital nonindependence of observations. We calculated relative risk for SMM by race/ethnicity using white women as the reference, within each study period.

We constructed an initial unadjusted model and a series of adjusted models. The initial unadjusted model included study period (baseline vs postintervention), race/ethnicity, and their interaction term. We then developed risk-adjusted models by adding maternal sociodemographic and clinical factors. We first adjusted for each covariate separately and compared the effect estimates between the unadjusted model and the single-covariate adjusted model. We then constructed a fully adjusted model by adding all covariates in the following sequence: (1) sociodemographic factors, (2) clinical factors except for the method of delivery, and (3) method of delivery. We added method of delivery separately from the other clinical factors to the model because all the other characteristics could also affect the delivery method and thus further influence SMM. Lastly, we performed sensitivity analysis by excluding each covariate one at a time from the fully adjusted model and evaluated the changes of the effect estimates. These analytical models were applied for both SMM and SMM excluding transfusion-only cases. All statistical analyses were performed using SAS 9.4 (SAS Institute Inc, Cary, NC).

gestational diabetes, previous cesarean delivery, labor induction, preterm birth), and method of delivery. All of these factors may contribute to racial inequalities in SMM among women with obstetric hemorrhage.

TABLE 2

Maternal characteristics of Hemorrhage Quality Collaborative Participants and California delivery population, baseline period: 2011–2014 (continued)

	Collaborative participants n (%)	All California n (%)
Prenatal care		
First trimester	825,239 (86.0)	1548609 (83.6)
Second trimester	106,875 (11.1)	239,552 (12.9)
Third trimester	23,143 (2.4)	54,128 (2.9)
No care	4,202 (0.4)	9,912 (0.5)
Missing	18,509	35,972
Chronic hypertension	17,012 (1.7)	31,024 (1.6)
Diabetes/gestational diabetes	106,227 (10.9)	187,960 (10.0)
Multiple gestation	17,946 (1.8)	31,058 (1.6)
Labor induction	150,786 (15.4)	286,710 (15.2)
Preterm birth (<37 wk gestation)	76,153 (7.8)	144,972 (7.7)
Previous cesarean delivery	165,297 (16.9)	331,037 (17.5)
Method of delivery: cesarean	316,141 (32.3)	622,465 (33.0)
Hospital characteristics		
Number of hospitals	99	238
Teaching hospitals	15 (15.2)	30 (12.6)
AAP neonatal level of care		
Level I	18 (18.2)	72 (30.3)
Level II	29 (29.3)	57 (23.9)
Level III	42 (42.4)	92 (38.7)
Level IV	10 (10.1)	17 (7.1)
Geographic region		
Central-South Coast	42 (42.4)	92 (38.7)
Central-North Coast and Northeastern	44 (44.4)	96 (40.3)
Central Valley, Southern Inland	13 (13.1)	50 (21.0)
Rural or urban-suburban		
Urban-Suburban	94 (94.9)	205 (86.1)
Rural	5 (5.1)	33 (13.9)
Average annual delivery volume (live births)		
<1000	15 (15.2)	72 (30.3)
1000–2499	55 (55.6)	117 (49.2)
≥3000	29 (29.3)	49 (20.6)
Hospital ownership		
University, city, county	11 (11.1)	39 (16.4)
Integrated health system	29 (29.3)	29 (12.2)
Private nonprofit	48 (48.5)	128 (53.8)
Private investor	11 (11.1)	42 (17.6)

AAP, American Academy of Pediatrics; BMI, body mass index; GED, General Educational Development.

Main et al. Reduction in racial disparities in severe maternal morbidity from hemorrhage in a large-scale quality improvement collaborative. *Am J Obstet Gynecol* 2020.

Results

Participating hospitals were diverse in size, ownership, neonatal intensive care level, the volume of deliveries, patient payer mix, and geography and were representative of the state as a whole (Table 2). Among the 54,311 women with obstetric hemorrhage at baseline, 42% were Hispanic, 29% were white, 15% were Asian, 5% were black, and 7% were other race/ethnicity. Racial/ethnic distribution in the postintervention period was similar to the baseline period. Maternal sociodemographic and clinical factors distributed differently across racial/ethnic groups (Table 3).

Overall reduction of SMM

Figure 1 shows the trend and the control chart of the quarterly SMM rate among all women with obstetric hemorrhage from 2011–2016. The total SMM rate in the baseline period was relatively stable (Figure 1, A). It dropped continuously after the initiation of the QI collaborative, and there was evidence of special cause of variation during the intervention and the postintervention period as illustrated by 4 consecutive points below the baseline lower control limit (3 standard deviations) and 8 consecutive points below the baseline average. The mean of the quarterly total SMM rate fell from 22.0% in the baseline to 18.6% in the postintervention period. The rate of SMM after excluding cases with transfusion alone also improved (Figure 1, B), with the rate reduced from 6.9% in the baseline to 5.6% in the postintervention period.

The reduction of SMM rate was observed in every racial/ethnic group, with absolute risk reductions ranged from 1.0% (20.5% baseline to 19.5% postintervention) in Asian women to 9.0% (28.6% baseline to 19.6% postintervention) in black women (Figure 2, A). Black women in the postintervention period were 23% less likely to have SMM (adjusted RR, 0.76; 95% CI, 0.65–0.89) as compared with those at baseline after adjusting for maternal sociodemographic and clinical factors. The rate of SMM after excluding transfusion-only cases also reduced in every racial/ethnic group (Figure 2, B). Although black

TABLE 3

Distribution of maternal sociodemographic and clinical factors among women with obstetric hemorrhage, by race/ethnicity and study period

Study period	Race/ethnicity											
	All women		White		Black		Asian		Hispanic		Other	
	Baseline	Postintervention	Baseline	Postintervention	Baseline	Postintervention	Baseline	Postintervention	Baseline	Postintervention	Baseline	Postintervention
Number of women with hemorrhage	54,311	19,165	15,775	5401	3404	1114	8180	3195	23,051	8161	3901	1294
Characteristics	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)
Age at delivery, y												
<18	906 (1.7)	200 (1.0)	76 (0.5)	15 (0.3)	91 (2.7)	14 (1.3)	13 (0.2)	4 (0.1)	702 (3.0)	164 (2.0)	24 (0.6)	3 (0.2)
18–25	12,774 (23.5)	3958 (20.7)	2700 (17.1)	707 (13.1)	1156 (34.0)	320 (28.7)	613 (7.5)	171 (5.4)	7737 (33.6)	2579 (31.6)	568 (14.6)	181 (14.0)
26–35	30,259 (55.7)	11,001 (57.4)	9674 (61.3)	3454 (64.0)	1622 (47.6)	557 (50.0)	5244 (64.1)	2085 (65.3)	11,391 (49.4)	4143 (50.8)	2328 (59.7)	762 (58.9)
≥36	10,372 (19.1)	4006 (20.9)	3325 (21.1)	1225 (22.7)	535 (15.7)	223 (20.0)	2310 (28.2)	935 (29.3)	3221 (14.0)	1275 (15.6)	981 (25.1)	348 (26.9)
Education												
Some high school or less	7486 (14.2)	2005 (10.8)	582 (3.8)	132 (2.5)	326 (10.0)	85 (8.0)	239 (3.0)	79 (2.6)	6203 (27.7)	1682 (21.4)	136 (3.6)	27 (2.1)
High school/GED	11,484 (21.8)	3956 (21.3)	2433 (15.8)	749 (14.3)	954 (29.1)	296 (27.8)	849 (10.8)	306 (9.9)	6732 (30.1)	2414 (30.7)	516 (13.6)	191 (15.2)
Some college	14,399 (27.3)	5123 (27.6)	4181 (27.2)	1341 (25.5)	1306 (39.9)	415 (39.0)	1399 (17.8)	489 (15.9)	6208 (27.7)	2443 (31.0)	1305 (34.5)	435 (34.5)
College grad or more	19,353 (36.7)	7446 (40.2)	8194 (53.2)	3027 (57.7)	687 (21.0)	269 (25.3)	5389 (68.4)	2211 (71.7)	3252 (14.5)	1332 (16.9)	1831 (48.3)	607 (48.2)
Missing	1589	635	385	152	131	49	304	110	656	290	113	34
Insurance												
Medi-Cal/other government-sponsored	18,697 (34.4)	6670 (34.8)	3079 (19.5)	1046 (19.4)	174 (49.2)	573 (51.4)	1246 (15.2)	532 (16.7)	11,949 (51.8)	4229 (51.8)	749 (19.2)	290 (22.4)
Private insurance	34,310 (63.2)	11,987 (62.5)	12,544 (79.5)	4301 (79.6)	1,673 (49.1)	528 (47.4)	6355 (77.7)	2376 (74.4)	10,630 (46.1)	3790 (46.4)	3108 (79.7)	992 (76.7)
Self-pay/no insurance	1303 (2.4)	508 (2.7)	151 (1.0)	54 (1.0)	57 (1.7)	13 (1.2)	579 (7.1)	287 (9.0)	472 (2.0)	142 (1.7)	44 (1.1)	12 (0.9)
Missing	1	0	1	0	0	0	0	0	0	0	0	0

Main et al. Reduction in racial disparities in severe maternal morbidity from hemorrhage in a large-scale quality improvement collaborative. Am J Obstet Gynecol 2020.

(continued)

TABLE 3

Distribution of maternal sociodemographic and clinical factors among women with obstetric hemorrhage, by race/ethnicity and study period (continued)

Study period	Race/ethnicity											
	All women		White		Black		Asian		Hispanic		Other	
	Baseline	Postintervention	Baseline	Postintervention	Baseline	Postintervention	Baseline	Postintervention	Baseline	Postintervention	Baseline	Postintervention
Number of women with hemorrhage	54,311	19,165	15,775	5401	3404	1114	8180	3195	23,051	8161	3901	1294
Characteristics	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)
Parity												
Multiparous	28,613 (52.7)	10,101 (52.7)	7260 (46.0)	2523 (46.7)	1972 (57.9)	668 (60.0)	3725 (45.5)	1498 (46.9)	13,470 (58.4)	4711 (57.7)	2186 (56.0)	701 (54.2)
Nulliparous	25,698 (47.3)	9064 (47.3)	8515 (54.0)	2878 (53.3)	1432 (42.1)	446 (40.0)	4455 (54.5)	1697 (53.1)	9581 (41.6)	3450 (42.3)	1715 (44.0)	593 (45.8)
Prepregnancy BMI												
<18.5	1079 (2.1)	372 (2.0)	297 (2.0)	93 (1.8)	73 (2.3)	18 (1.7)	356 (4.8)	156 (5.1)	286 (1.3)	88 (1.1)	67 (1.9)	17 (1.4)
18.5–25	22,140 (43.8)	7597 (41.1)	7692 (51.7)	2570 (48.9)	1015 (32.4)	334 (31.0)	4722 (64.3)	1919 (62.6)	7051 (32.7)	2287 (29.1)	1660 (45.9)	487 (39.1)
25–30	14,347 (28.4)	5336 (28.9)	3837 (25.8)	1442 (27.5)	887 (28.3)	295 (27.3)	1636 (22.3)	703 (22.9)	6919 (32.1)	2500 (31.9)	1068 (29.5)	396 (31.8)
30–35	7485 (14.8)	2953 (16.0)	1689 (11.4)	660 (12.6)	573 (18.3)	190 (17.6)	477 (6.5)	218 (7.1)	4267 (19.8)	1691 (21.5)	479 (13.2)	194 (15.6)
35–40	3287 (6.5)	1317 (7.1)	791 (5.3)	283 (5.4)	290 (9.3)	125 (11.6)	116 (1.6)	55 (1.8)	1872 (8.7)	764 (9.7)	218 (6.0)	90 (7.2)
≥40	2202 (4.4)	914 (4.9)	559 (3.8)	205 (3.9)	297 (9.5)	117 (10.8)	38 (0.5)	13 (0.4)	1182 (5.5)	517 (6.6)	126 (3.5)	62 (5.0)
Missing	3,771	676	910	148	269	35	835	131	1,474	314	283	48
Prenatal care												
First trimester	45,997 (86.2)	16,391 (86.9)	13,854 (89.5)	4778 (89.9)	2733 (82.6)	923 (85.1)	7245 (90.7)	2839 (90.0)	18,871 (83.1)	6728 (83.7)	3294 (85.3)	1123 (87.7)
Second trimester	5781 (10.8)	1902 (10.1)	1262 (8.2)	407 (7.7)	422 (12.8)	123 (11.3)	622 (7.8)	254 (8.0)	3022 (13.3)	994 (12.4)	453 (11.7)	124 (9.7)
Third trimester	1192 (2.2)	435 (2.3)	249 (1.6)	91 (1.7)	103 (3.1)	30 (2.8)	105 (1.3)	61 (1.9)	642 (2.8)	233 (2.9)	93 (2.4)	20 (1.6)
No care	381 (0.7)	143 (0.8)	114 (0.7)	37 (0.7)	50 (1.5)	9 (0.8)	18 (0.2)	2 (0.1)	176 (0.8)	82 (1.0)	23 (0.6)	13 (1.0)
Missing	960	294	296	88	96	29	190	39	340	124	38	14

Main et al. Reduction in racial disparities in severe maternal morbidity from hemorrhage in a large-scale quality improvement collaborative. Am J Obstet Gynecol 2020.

(continued)

TABLE 3

Distribution of maternal sociodemographic and clinical factors among women with obstetric hemorrhage, by race/ethnicity and study period (continued)

Study period	Race/ethnicity											
	All women		White		Black		Asian		Hispanic		Other	
	Baseline	Postintervention	Baseline	Postintervention	Baseline	Postintervention	Baseline	Postintervention	Baseline	Postintervention	Baseline	Postintervention
Number of women with hemorrhage	54,311	19,165	15,775	5401	3404	1114	8180	3195	23,051	8161	3901	1294
Characteristics	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)
Chronic hypertension												
No	52,981 (97.6)	18,389 (96.0)	15,424 (97.8)	5215 (96.6)	3218 (94.5)	1025 (92.0)	8046 (98.4)	3118 (97.6)	22,555 (97.8)	7845 (96.1)	3738 (95.8)	1186 (91.7)
Yes	1330 (2.4)	776 (4.0)	351 (2.2)	186 (3.4)	186 (5.5)	89 (8.0)	134 (1.6)	77 (2.4)	496 (2.2)	316 (3.9)	163 (4.2)	108 (8.3)
Gestational diabetes												
No	47,210 (86.9)	16,511 (86.2)	14,449 (91.6)	4928 (91.2)	3047 (89.5)	985 (88.4)	6802 (83.2)	2626 (82.2)	19,885 (86.3)	6962 (85.3)	3027 (77.6)	1010 (78.1)
Yes	7101 (13.1)	2654 (13.8)	1326 (8.4)	473 (8.8)	357 (10.5)	129 (11.6)	1378 (16.8)	569 (17.8)	3166 (13.7)	1199 (14.7)	874 (22.4)	284 (21.9)
Multiple gestation												
No	51,974 (95.7)	18,377 (95.9)	14,847 (94.1)	5108 (94.6)	3246 (95.4)	1075 (96.5)	7798 (95.3)	3042 (95.2)	22,290 (96.7)	7904 (96.9)	3793 (97.2)	1248 (96.4)
Yes	2337 (4.3)	788 (4.1)	928 (5.9)	293 (5.4)	158 (4.6)	39 (3.5)	382 (4.7)	153 (4.8)	761 (3.3)	257 (3.1)	108 (2.8)	46 (3.6)
Labor induction												
No	44,015 (81.0)	15,291 (79.8)	12,385 (78.5)	4184 (77.5)	2736 (80.4)	888 (79.7)	6703 (81.9)	2595 (81.2)	19,044 (82.6)	6559 (80.4)	3147 (80.7)	1065 (82.3)
Yes	10,296 (19.0)	3874 (20.2)	3390 (21.5)	1,217 (22.5)	668 (19.6)	226 (20.3)	1477 (18.1)	600 (18.8)	4007 (17.4)	1602 (19.6)	754 (19.3)	229 (17.7)
Preterm birth												
No	44,077 (81.2)	15,803 (82.5)	12,855 (81.5)	4461 (82.6)	2509 (73.8)	851 (76.5)	6832 (83.6)	2718 (85.1)	18,709 (81.2)	6711 (82.3)	3172 (81.4)	1062 (82.1)
Yes	10,206 (18.8)	3352 (17.5)	2914 (18.5)	937 (17.4)	889 (26.2)	262 (23.5)	1344 (16.4)	476 (14.9)	4334 (18.8)	1445 (17.7)	725 (18.6)	232 (17.9)
Missing	28	10	6	3	6	1	4	1	8	5	4	0

Main et al. Reduction in racial disparities in severe maternal morbidity from hemorrhage in a large-scale quality improvement collaborative. Am J Obstet Gynecol 2020.

(continued)

TABLE 3
Distribution of maternal sociodemographic and clinical factors among women with obstetric hemorrhage, by race/ethnicity and study period (continued)

Study period		Race/ethnicity											
		All women			White			Black			Asian		
		Baseline			Postintervention			Baseline			Postintervention		
		n (%)			n (%)			n (%)			n (%)		
Number of women with hemorrhage		54,311			15,775			3404			8180		
Characteristics		n (%)			n (%)			n (%)			n (%)		
Previous cesarean delivery													
No		46,179 (85.0)			13,843 (87.8)			2719 (79.9)			7045 (86.1)		
Yes		8132 (15.0)			1932 (12.2)			685 (20.1)			1135 (13.9)		
Method of delivery													
Vaginal		32,717 (60.2)			9798 (62.1)			1729 (50.8)			4702 (57.5)		
Cesarean		21,594 (39.8)			5977 (37.9)			1675 (49.2)			3478 (42.5)		

BMI, body mass index; GED, General Educational Development.

Main et al. Reduction in racial disparities in severe maternal morbidity from hemorrhage in a large-scale quality improvement collaborative. Am J Obstet Gynecol 2020.

women experienced the largest reduction (2.5% absolute rate reduction), their rate was still the greatest among all racial/ethnic groups in the post-intervention period (6.9% vs 5.1–5.5%).

Risk adjustment models for comparing racial differences


We then assessed racial differences in the risk of SMM among women with obstetric hemorrhage, using white women as the reference. At baseline, black mothers were associated with a greater risk of SMM compared with white mothers (RR, 1.34; 95% CI, 1.26–1.42) (Table 4, model 1). The baseline racial difference was significant even after adjusting for all of the sociodemographic and clinical factors (RR, 1.22; 95% CI, 1.15–1.30) (Table 4, model 4). In the postintervention period, the black–white RR decreased to 1.22 in the unadjusted model, although it remained significant (95% CI, 1.05–1.40) (Table 4, model 1). However, once clinical factors and especially method of delivery (eg, cesarean) were added to the adjustment model, the racial difference were no longer significant (RR, 1.14; 95% CI, 0.98–1.32) (Table 4, model 3), and (RR, 1.07; 95% CI, 0.92–1.24) (Table 4, model 4). Among all the covariates, method of delivery influenced the black–white relative risk the most (Supplemental Tables 2 and 3).

Results for SMM excluding transfusion-only cases were similar. In this group of women, the racial/ethnic inequality between black and white women for SMM decreased from baseline (RR, 1.46; 95% CI, 1.30–1.65) to the postintervention period (RR, 1.22; 95% CI, 0.98–1.52, (Table 3, model 1). After adjusting for all of the covariates, the significant black–white differences at baseline (RR, 1.33; 95% CI, 1.16–1.52) were attenuated in the postintervention period (RR, 0.99; 95% CI, 0.76–1.28, (Table 4, model 4). Again, results from the single-covariate model and sensitivity analysis suggested that the method of delivery was the most influential covariate (Supplemental Tables 4 and 5).


FIGURE 1

Control chart of quarterly rates for SMM, 2011–2016

A Severe maternal morbidity



B Severe maternal morbidity excluding cases with transfusion alone



A, SMM. B, SMM excluding cases with transfusion alone, 2011–2016.

SMM, severe maternal morbidity.

Main et al. Reduction in racial disparities in severe maternal morbidity from hemorrhage in a large-scale quality improvement collaborative. *Am J Obstet Gynecol* 2020.

Sensitivity analyses

Because some of the SMM diagnoses may not be directly related to excessive bleeding, we performed additional sensitivity analysis restricting our analyses to SMM diagnoses that had the

strongest relationship to hemorrhage, including transfusion; acute renal failure; adult respiratory distress; cardiac arrest; disseminated intravascular coagulation; acute heart failure; pulmonary edema; shock; hysterectomy;

and ventilation. Almost 99% of SMM cases and almost 93% of SMM cases excluding transfusion alone were related to hemorrhage, and all results were nearly identical to those presented (Supplemental Tables 6 and 7).

Comment



Large variations in rates of SMM exist across hospitals in the United States.¹ Both “within-hospital” and “between-hospital” disparities in SMM have been documented,¹³ and variations in the quality of care delivery have been shown to contribute to racial/ethnic disparities in SMM.²⁸ Obstetric hemorrhage is an acute delivery event that accounts for approximately one half of SMM. Importantly, maternal deaths and severe complications from hemorrhage have been judged to have a high degree of preventability largely due to provider improvement opportunities.²⁹ Therefore, reducing variations in clinical care processes by implementing standard protocols has the potential to both improve hemorrhage outcomes and concurrently reduce racial/ethnic disparities. Obstetric hemorrhage provides a useful model to explore the impact of QI efforts on reducing disparities in maternal outcomes.

Principal findings

In this cross-sectional study among women with obstetric hemorrhage in the 99 hospitals participated in the QI collaborative, we showed that SMM rate was reduced in every racial/ethnic group after the intervention. Of particular notice, the risk of SMM was no longer greater in black women compared with white women in the postintervention period after accounting for sociodemographic and clinical factors. The marked improvement in black rates of SMM from hemorrhage and the narrowing of black–white difference are important findings suggesting QI efforts can be effective in both improving maternal outcomes and reducing inequities in care delivery for a specific medical condition.

Clinical implications

Results from adjusted analyses indicated that the remaining racial inequalities in SMM could be largely

FIGURE 2
Rates of SMM**A Severe maternal morbidity****B Severe maternal morbidity excluding cases with transfusion alone**

Rate of **A**, SMM and **B**, SMM excluding cases with transfusion alone, by race/ethnicity and study period. Poisson generalized estimating equation model included study period, race/ethnicity, the interaction term between study period and race/ethnicity, and all of the maternal sociodemographic and clinical factors listed in Table 4.

adjRR, adjusted relative risk; CI, confidence interval. N, Number of denominator (women with obstetric hemorrhage); SMM, severe maternal morbidity.

Main et al. Reduction in racial disparities in severe maternal morbidity from hemorrhage in a large-scale quality improvement collaborative. Am J Obstet Gynecol 2020.

explained by controlled sociodemographic and clinical factors with the method of delivery being the most influential factor. Cesarean delivery has been estimated to be associated with a larger proportion of SMM than was any other risk factor.³⁰ Currently, black women have greater rates of cesarean delivery compared with women in other race/ethnic groups, but this difference only began in the 1990s.³¹ Reducing disparity in the cesarean rate, where possible, may allow further reduction in black–white difference in SMM from hemorrhage.

Blood transfusions are an important driver for SMM cases. The greater transfusion rate among black mothers may be partially related to greater rates of anemia among black women when presenting for delivery (2–6 times greater than white women).^{32–34} Anemia, in turn, increases the risk of transfusion, particularly in the setting of surgical delivery. Improved recognition and treatment of anemia before delivery may be another approach to reduce black–white disparity in SMM.

The subset of women with SMM excluding transfusion-only cases represents a group of women with potentially more severe diagnoses or procedures including hysterectomy, renal or respiratory failure. Among these women, a more pronounced narrowing of black–white difference was observed. Again, greater cesarean delivery rates among black women appeared to account for a large portion of the remaining difference in this group of women.

Research implications

QI has been suggested as an important approach to address racial/ethnic disparities.³⁵ However, in practice, QI efforts have been shown to have a variable impact on racial/ethnic disparities in outcomes.^{20–23,36} There is limited experience examining the potential for QI to reduce obstetric disparities. Our study provides several insights to better understand the application of QI for successful reduction of disparities.

TABLE 4

Association between race/ethnicity and SMM, both total and excluding cases with transfusion alone, by study period

Race/ethnicity	Total SMM		SMM excluding cases with transfusion alone ^a	
	Baseline	Postintervention	Baseline	Postintervention
	RR (95% CI)	RR (95% CI)	RR (95% CI)	RR (95% CI)
Model 1 ^b : Unadjusted model				
White	1.00	1.00	1.00	1.00
Black	1.34 (1.27–1.42)	1.22 (1.05–1.40)	1.46 (1.30–1.65)	1.22 (0.98–1.52)
Asian	1.03 (0.95–1.11)	1.13 (1.00–1.28)	1.15 (1.05–1.25)	1.09 (0.90–1.32)
Hispanics	1.09 (1.03–1.15)	1.10 (0.99–1.23)	1.05 (0.95–1.15)	0.96 (0.82–1.12)
Others	1.08 (1.00–1.18)	1.07 (0.92–1.25)	1.02 (0.87–1.19)	0.97 (0.75–1.27)
Model 2 ^c : Adjusted for sociodemographic factors				
White	1.00	1.00	1.00	1.00
Black	1.32 (1.25–1.39)	1.18 (1.02–1.37)	1.47 (1.30–1.66)	1.16 (0.91–1.49)
Asian	1.02 (0.96–1.09)	1.12 (0.98–1.26)	1.11 (1.01–1.21)	1.05 (0.86–1.27)
Hispanics	1.07 (1.01–1.12)	1.06 (0.95–1.18)	1.04 (0.95–1.13)	0.94 (0.80–1.12)
Others	1.07 (0.98–1.17)	1.06 (0.90–1.23)	0.99 (0.84–1.16)	0.96 (0.73–1.26)
Model 3 ^d : Adjusted for sociodemographic and clinical factors (except for method of delivery)				
White	1.00	1.00	1.00	1.00
Black	1.29 (1.21–1.38)	1.14 (0.98–1.32)	1.43 (1.25–1.63)	1.05 (0.81–1.37)
Asian	1.03 (0.97–1.10)	1.09 (0.97–1.23)	1.09 (0.99–1.20)	1.05 (0.86–1.27)
Hispanics	1.09 (1.04–1.15)	1.07 (0.97–1.17)	1.05 (0.96–1.16)	0.96 (0.83–1.13)
Others	1.09 (1.00–1.19)	1.04 (0.89–1.21)	0.98 (0.83–1.16)	0.94 (0.72–1.23)
Model 4 ^e : Adjusted for sociodemographic, and clinical factors including method of delivery (fully adjusted model)				
White	1.00	1.00	1.00	1.00
Black	1.22 (1.15–1.30)	1.07 (0.92–1.24)	1.33 (1.16–1.52)	0.99 (0.76–1.29)
Asian	1.02 (0.96–1.09)	1.09 (0.96–1.23)	1.06 (0.96–1.18)	1.05 (0.86–1.27)
Hispanics	1.09 (1.04–1.15)	1.06 (0.97–1.16)	1.05 (0.95–1.16)	0.96 (0.82–1.12)
Others	1.07 (0.98–1.16)	1.03 (0.89–1.21)	0.95 (0.80–1.12)	0.92 (0.70–1.22)

CI, confidence interval; RR, relative risk; SMM, severe maternal morbidity.

^a Cases with transfusion alone were excluded from the numerator; ^b Unadjusted model. Model included study period, race ethnicity, and the interaction term between study period and race ethnicity;^c Based on model 1, additionally included the following maternal sociodemographic factors: age at delivery, education levels, parity, and insurance status; ^d Based on model 2, additionally included the following maternal clinical factors: prenatal visits, pre-pregnancy body mass index, multiple pregnancy, chronic hypertension, gestational diabetes, previous cesarean delivery, labor induction, and pre-term birth; ^e Fully adjusted model. Based on model 3, additionally included method of delivery.

Main et al. Reduction in racial disparities in severe maternal morbidity from hemorrhage in a large-scale quality improvement collaborative. Am J Obstet Gynecol 2020.

First, QI efforts may have variable effectiveness because the focus of the intervention does not sufficiently address the primary sources of racial/ethnic disparities. Social determinants, underlying risk factors, structural racism, lack of trust and respect from providers, and the quality of the care received all contribute to disparities in maternal outcomes.¹³ In addition, specific causes of mortality and morbidity may be more or less amenable to QI

efforts. Evaluating the relative contribution of these factors to performance gaps among racial/ethnic groups is important in designing effective interventions. We suggest, similar to Wise,³⁷ that QI efforts for reducing disparities are the most likely to be successful when they (1) target care-sensitive conditions (2) for an acute process (3) where poor access to highly effective treatment (4) is the dominant reason for the morbidity. Management

of obstetric hemorrhage offers such an example. Case reviews of SMM due to hemorrhage have consistently identified provider and system-related factors as the dominant contributors to adverse outcomes.^{8,38} At the same time, implementation of systematic approaches to management of hemorrhage have been shown to reduce rates of SMM.^{18,39} For other conditions that are less acute, more closely tied to patient risk factors and chronic

exposure to racism, solutions will need to be more broadly based. We suggest that eliminating racial disparities will require appropriate matching between the source of disparity, the causal pathway and the intervention(s).

Limitations and strengths

The limitations of our study must be viewed in light of its design. As we were looking for population effects, we relied on administrative data for outcomes. However, we took extra steps for validation of outcomes including case reviews and outlier checks. The risk of coding variation was minimized by following the same hospitals with the same racial mix over time. Although we included a broad range of maternal risk factors associated with SMM in the adjusted model, we may not have accounted for unmeasured confounders. In addition, even with our very large number of mothers, we may have limited sample size for detecting statistical interactions.

Our study has several strengths. Self-reported racial/ethnic data were obtained from birth certificates, which are generally regarded as the gold standard. Our intervention reached a large sample of women and included a broad range of hospital sizes and affiliations that collectively care for more than 250,000 births annually. We focused on a defined subset of women diagnosed with hemorrhage that allowed us to deploy a clearly defined intervention bundle with nationally recognized outcome measures.

Conclusion

Our findings point to opportunities for reduction in black–white disparity for the most common maternity complication, hemorrhage, by implementing national safety bundles for prevention and response to obstetric hemorrhage. In addition, our data suggest that efforts to decrease the greater rate of cesarean deliveries among black women may show added benefit. The greater rate of anemia at labor admission in black women also provides a prenatal care improvement opportunity that could impact transfusion rates. These clinical efforts should be in parallel with efforts to reverse bias

and racism in the medical system by treating black women with respect and dignity, better understanding their circumstances, and listening to and acting on their concerns. All of these approaches are necessary to address the persistent disparities that we see in obstetric care. ■

Acknowledgments

We wish to acknowledge the 40 mentors and the many hospital teams who led this effort at the facility level.

References

1. Callaghan WM. Overview of maternal mortality in the United States. *Semin Perinatol* 2012;36:2–6.
2. Creanga AA, Berg CJ, Syverson C, Seed K, Bruce FC, Callaghan WM. Race, ethnicity, and nativity differentials in pregnancy-related mortality in the United States: 1993–2006. *Obstet Gynecol* 2012;120:261–8.
3. Grobman WA, Bailit JL, Rice MM, et al. Racial and ethnic disparities in maternal morbidity and obstetric care. *Obstet Gynecol* 2015;125:1460–7.
4. Centers for Disease Control and Prevention. Severe Maternal Morbidity Indicators and Corresponding ICD Codes during Delivery Hospitalizations. Available at: <https://www.cdc.gov/reproductivehealth/maternalinfanthealth/smm/severe-morbidity-ICD.htm>. Accessed December 6, 2018.
5. Callaghan WM, Creanga AA, Kuklina EV. Severe maternal morbidity among delivery and postpartum hospitalizations in the United States. *Obstet Gynecol* 2012;120:1029–36.
6. Creanga AA, Berg CJ, Ko JY, et al. Maternal mortality and morbidity in the United States: where are we now? *J Womens Health (Larchmt)* 2014;23:3–9.
7. Creanga AA, Bateman BT, Kuklina EV, Callaghan WM. Racial and ethnic disparities in severe maternal morbidity: a multistate analysis, 2008–2010. *Am J Obstet Gynecol* 2014;210:435.e1–8.
8. Berg CJ, Harper MA, Atkinson SM, et al. Preventability of pregnancy-related deaths: results of a state-wide review. *Obstet Gynecol* 2005;106:1228–34.
9. Berg CJ, Mackay AP, Qin C, Callaghan WM. Overview of maternal morbidity during hospitalization for labor and delivery in the United States: 1993–1997 and 2001–2005. *Obstet Gynecol* 2009;113:1075–81.
10. Danel I, Berg C, Johnson CH, Atrash H. Magnitude of maternal morbidity during labor and delivery: United States, 1993–1997. *Am J Public Health* 2003;93:631–4.
11. Geller SE, Cox SM, Callaghan WM, Berg CJ. Morbidity and mortality in pregnancy: laying the groundwork for safe motherhood. *Womens Health Issues* 2006;16:176–88.
12. Howell EA, Egorova NN, Balbierz A, Zeitlin J, Hebert PL. Site of delivery contribution to black-white severe maternal morbidity disparity. *Am J Obstet Gynecol* 2016;215:143–52.
13. Howell EA, Zeitlin J. Improving hospital quality to reduce disparities in severe maternal morbidity and mortality. *Semin Perinatol* 2017;41:266–72.
14. Bateman BT, Berman MF, Riley LE, Leffert LR. The epidemiology of postpartum hemorrhage in a large, nationwide sample of deliveries. *Anesth Analg* 2010;110:1368–73.
15. Callaghan WM, Kuklina EV, Berg CJ. Trends in postpartum hemorrhage: United States, 1994–2006. *Am J Obstet Gynecol* 2010;202:353.e1–6.
16. Khan KS, Wojdyla D, Say L, Gulmezoglu AM, Van Look PF. WHO analysis of causes of maternal death: a systematic review. *Lancet* 2006;367:1066–74.
17. Kramer MS, Berg C, Abenhaim H, et al. Incidence, risk factors, and temporal trends in severe postpartum hemorrhage. *Am J Obstet Gynecol* 2013;209:449.e1–7.
18. Main EK, Goffman D, Scavone BM, et al. National Partnership for Maternal Safety: consensus bundle on obstetric hemorrhage. *Obstet Gynecol* 2015;126:155–62.
19. Weinick RM, Hasnain-Wynia R. Quality improvement efforts under health reform: how to ensure that they help reduce disparities—not increase them. *Health Aff (Millwood)* 2011;30:1837–43.
20. Areal PA, Ayalon L, Hunkeler E, et al. Improving depression care for older, minority patients in primary care. *Med Care* 2005;43:381–90.
21. Sequist TD, Adams A, Zhang F, Ross-Degnan D, Ayanian JZ. Effect of quality improvement on racial disparities in diabetes care. *Arch Intern Med* 2006;166:675–81.
22. Trivedi AN, Grebla RC, Wright SM, Washington DL. Despite improved quality of care in the Veterans Affairs health system, racial disparity persists for important clinical outcomes. *Health Aff (Millwood)* 2011;30:707–15.
23. Trivedi AN, Zaslavsky AM, Schneider EC, Ayanian JZ. Trends in the quality of care and racial disparities in Medicare managed care. *N Engl J Med* 2005;353:692–700.
24. Werner RM, Asch DA, Polsky D. Racial profiling: the unintended consequences of coronary artery bypass graft report cards. *Circulation* 2005;111:1257–63.
25. Main EK, Cape V, Abreo A, et al. Reduction of severe maternal morbidity from hemorrhage using a state perinatal quality collaborative. *Am J Obstet Gynecol* 2017;216:298.e1–11.
26. Main EK, Dhurjati R, Cape V, et al. Improving maternal safety at scale with the mentor model of collaborative improvement. *Jt Comm J Qual Patient Saf* 2018;44:250–9.

27. Beate D. Probabilistic record linkages for generating a comprehensive epidemiological research file on maternal and infant health. Available at: <http://www.health-info-solutions.com/Probabilistic%20Record%20Linkages%20may%202002.pdf>. Accessed September 1, 2018.
28. Howell EA, Egorova N, Balbierz A, Zeitlin J, Hebert PL. Black-white differences in severe maternal morbidity and site of care. *Am J Obstet Gynecol* 2016;214:122.e1–7.
29. Main EK, McCain CL, Morton CH, Holtby S, Lawton ES. Pregnancy-related mortality in California: causes, characteristics, and improvement opportunities. *Obstet Gynecol* 2015;125:938–47.
30. Leonard SA, Main EK, Carmichael SL. The contribution of maternal characteristics and cesarean delivery to an increasing trend of severe maternal morbidity. *BMC Pregnancy Childbirth* 2019;19:16.
31. Taffel SM. Cesarean delivery in the United States, 1990. *Vital and health statistics Series 21, Data on natality, marriage, and divorce*. Hyattsville, MD: US Department of Health and Human Services; 1994:1–24.
32. Adebisi OY, Strayhorn G. Anemia in pregnancy and race in the United States: blacks at risk. *Fam Med* 2005;37:655–62.
33. Le CH. The prevalence of anemia and moderate-severe anemia in the US population (NHANES 2003-2012). *PLoS One* 2016;11:e0166635.
34. Mohamed MA, Ahmad T, Macri C, Aly H. Racial disparities in maternal hemoglobin concentrations and pregnancy outcomes. *J Perinat Med* 2012;40:141–9.
35. Aaron KF, Clancy CM. Improving quality and reducing disparities: toward a common pathway. *JAMA* 2003;289:1033–4.
36. Sehgal AR. Impact of quality improvement efforts on race and sex disparities in hemodialysis. *JAMA* 2003;289:996–1000.
37. Wise PH. The anatomy of a disparity in infant mortality. *Ann Rev Public Health* 2003;24:341–62.
38. Geller SE, Rosenberg D, Cox SM, et al. The continuum of maternal morbidity and mortality: factors associated with severity. *Am J Obstet Gynecol* 2004;191:939–44.
39. Shields LE, Wiesner S, Fulton J, Pelletreau B. Comprehensive maternal hemorrhage protocols reduce the use of blood products and improve patient safety. *Am J Obstet Gynecol* 2015;212:272–80.

Author and article information

From the Division of Maternal-Fetal Medicine, Department of Obstetrics and Gynecology (Dr Main) and Division

of Neonatal and Developmental Medicine, Department of Pediatrics (Drs Chang, Dhurjati, Cape, Profit, and Gould), Stanford University School of Medicine; and California Maternal Quality Care Collaborative (Drs Main, Chang, Dhurjati, Cape, and Gould) and California Perinatal Quality Care Collaborative (Drs Dhurjati, Profit, and Gould), Stanford, CA.

Received Sept. 3, 2019; revised Dec. 18, 2019; accepted Jan. 13, 2020.

The authors report no conflict of interest.

The research in this publication was supported by funding from Merck, through its Merck for Mothers program, and is the sole responsibility of the authors. Merck for Mothers is known as MSD for Mothers outside the United States and Canada. The funder has no involvement in study design, collection, analysis, or interpretation of results. J.P.'s effort was supported in part by grants from the *Eunice Kennedy Shriver* National Institute of Child Health and Human Development (R01 HD083368-01 and R01 HD08467-01, PI J.P.). The content is solely the responsibility of the authors and does not necessarily represent the official views of the *Eunice Kennedy Shriver* National Institute of Child Health and Human Development or the National Institutes of Health. This funding source had no role in any of the following aspects of this study: design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

Corresponding author: Elliott K. Main, MD. emain@stanford.edu