

Safe Motherhood Initiative



**ACOG**  
THE AMERICAN CONGRESS  
OF OBSTETRICIANS  
AND GYNECOLOGISTS

District II

# Maternal Safety Bundle for Severe Hypertension in Pregnancy

REVISED NOVEMBER 2015

Disclaimer: The following material is an example only and not meant to be prescriptive. ACOG accepts no liability for the content or for the consequences of any actions taken on the basis of the information provided.

# KEY ELEMENTS

## RISK ASSESSMENT & PREVENTION

- Diagnostic Criteria
- When to Treat
- Agents to Use
- Monitoring

## READINESS & RESPONSE

- Complications & Escalation Process
- Further Evaluation
- Change of Status
- Postpartum Surveillance

# TYPES OF HYPERTENSION

EXAMPLE

4

CHRONIC HYPERTENSION	<ul style="list-style-type: none"> <li>○ SBP <math>\geq</math> 140 <b>or</b> DBP <math>\geq</math> 90</li> <li>○ Pre-pregnancy or <math>&lt;</math>20 weeks</li> </ul>
GESTATIONAL HYPERTENSION	<ul style="list-style-type: none"> <li>○ SBP <math>\geq</math> 140 <b>or</b> DBP <math>\geq</math> 90</li> <li>○ <math>&gt;</math> 20 weeks</li> <li>○ Absence of proteinuria or systemic signs/symptoms</li> </ul>
PREECLAMPSIA - ECLAMPSIA	<ul style="list-style-type: none"> <li>○ SBP <math>\geq</math> 140 <b>or</b> DBP <math>\geq</math> 90</li> <li>○ Proteinuria with or without signs/symptoms</li> <li>○ Presentation of signs/symptoms/lab abnormalities but no proteinuria</li> </ul> <p><i>*Proteinuria not required for diagnosis eclampsia seizure in setting of preeclampsia</i></p>
CHRONIC HYPERTENSION + SUPERIMPOSED PREECLAMPSIA	
<p><b>PREECLAMPSIA WITH SEVERE FEATURES</b></p>	<ul style="list-style-type: none"> <li>○ Two severe BP values (SBP <math>\geq</math> 160 <b>or</b> DBP <math>\geq</math> 110) obtained 15-60 minutes apart</li> <li>○ Persistent oliguria <math>&lt;</math>500 ml/24 hours</li> <li>○ Progressive renal insufficiency</li> <li>○ Unremitting headache/visual disturbances</li> <li>○ Pulmonary edema</li> <li>○ Epigastric/RUQ pain</li> <li>○ LFTs <math>&gt;</math> 2x normal</li> <li>○ Platelets <math>&lt;</math> 100K</li> <li>○ HELLP syndrome</li> </ul> <p><i>*5 gr of proteinuria no longer criteria for severe preeclampsia</i></p>

# DEFINITIONS

## SEVERE HYPERTENSION

---

- Systolic blood pressure  $\geq$  160 mm Hg or
- Diastolic blood pressure  $\geq$  110 mm Hg

## HYPERTENSIVE EMERGENCY

---

- Persistent, severe hypertension that can occur antepartum, intrapartum, or postpartum
- Defined as:
  - Two severe BP values ( $\geq$  160/110) taken 15-60 minutes apart
  - Severe values do not need to be consecutive

# WHEN TO TREAT

EXAMPLE

6

## SEVERE HYPERTENSION

SBP  $\geq$  160 **or** DBP  $\geq$  110

- Repeat BP every 5 min for 15 min
- Notify physician after one severe BP value is obtained

## HYPERTENSIVE EMERGENCY

Persistent, severe hypertension that can occur antepartum, intrapartum, or postpartum

Two severe BP values ( $\geq$  160/110) taken 15-60 minutes apart

Severe values do not need to be consecutive

- If severe BP elevations persist for 15 min or more, begin treatment **ASAP. Preferably within 60 min of the second elevated value.**
- If two severe BPs are obtained *within* 15 min, treatment may be initiated if clinically indicated

# FIRST LINE THERAPIES

EXAMPLE

7

- Intravenous labetalol
- Intravenous hydralazine
- Oral nifedipine

## Magnesium sulfate not recommended as antihypertensive agent

- Should be used for: seizure prophylaxis and controlling seizures in eclampsia
- IV bolus of 4-6 grams in 100 ml over 20 minutes, followed by IV infusion of 1-2 grams per hour. **Continue for 24 hours postpartum**
- If no IV access, 10 grams of 50% solution IM (5 g in each buttock)
- Contraindications: pulmonary edema, renal failure, myasthenia gravis

## Anticonvulsants (for recurrent seizures or when magnesium is C/I):

- **Lorazepam:** 2-4 mg IV x 1, may repeat x 1 after 10-15 min
- **Diazepam:** 5-10 mg IV every 5-10 min to max dose 30 mg
- **Phenytoin:** 15-20 mg/kg IV x 1, may repeat 10 mg/kg IV after 20 min if no response. Avoid with hypotension, may cause cardiac arrhythmias.
- **Keppra:** 500 mg IV or orally, may repeat in 12 hours. Dose adjustment needed if renal impairment.

*\*There may be adverse effects and additional contraindications. Clinical judgement should prevail*

# Labetalol Algorithm

## EXAMPLE

Trigger: If severe elevations (SBP  $\geq 160$  or DBP  $\geq 110$ ) persist for 15 min or more **OR** If two severe elevations are obtained within 15 min and tx is clinically indicated



- Notify provider after one severe BP value is obtained
- Institute fetal surveillance if viable
- Hold IV labetalol for maternal pulse under 60
- Maximum cumulative IV-administered dose of labetalol should not exceed 220 mg in 24 hours
- There may be adverse effects and contraindications. Clinical judgement should prevail.

# Hydralazine Algorithm

EXAMPLE

Trigger: If severe elevations (SBP  $\geq 160$  or DBP  $\geq 110$ ) persist for 15 min or more **OR** If two severe elevations are obtained within 15 min and tx is clinically indicated




- Notify provider after one severe BP value is obtained
- Institute fetal surveillance if viable
- Hold IV labetalol for maternal pulse under 60
- Maximum cumulative IV-administered dose of hydralazine should not exceed 25 mg in 24 hours
- There may be adverse effects and contraindications. Clinical judgement should prevail.

# Oral Nifedipine Algorithm

EXAMPLE

Trigger: If severe elevations (SBP  $\geq 160$  or DBP  $\geq 110$ ) persist for 15 min or more **OR** If two severe elevations are obtained within 15 min and tx is clinically indicated



- Notify provider after one severe BP value is obtained
- Institute fetal surveillance if viable
- Capsules should be administered orally and not punctured or otherwise administered sublingually
- There may be adverse effects and contraindications. Clinical judgement should prevail.

# ADDITIONAL THERAPY RECOMMENDATIONS

## IF NO IV ACCESS AVAILABLE:

- Initiate algorithm for oral nifedipine, or
- Oral labetalol, 200 mg *\*Repeat in 30 min if SBP remains  $\geq 160$  or DBP  $\geq 110$  and IV access still unavailable*

## SECOND LINE THERAPIES (if patient fails to respond to first line tx):

Recommend emergency consult with:

- Maternal Fetal Medicine
- Internal Medicine
- Anesthesiology
- Critical Care
- Emergency Medicine

*May also consider:*

- ✓ Labetalol or nicardipine via infusion pump
- ✓ Sodium nitroprusside for extreme emergencies *\*Use for shortest amount of time due to cyanide/thiocyanate toxicity*

# MONITORING BLOOD PRESSURE

## MATERNAL

- Once BP is controlled (<160/110), measure
  - ✓ Every 10 minutes for 1 hour
  - ✓ Every 15 minutes for next hour
  - ✓ Every 30 minutes for next hour
  - ✓ Every hour for 4 hours
- Obtain baseline labs:
  - ✓ CBC
  - ✓ Platelets
  - ✓ LDH
  - ✓ Liver Function Tests
  - ✓ Electrolytes
  - ✓ BUN creatinine
  - ✓ Urine protein

## FETAL

- Fetal monitoring surveillance as appropriate for gestational age

# Hypertensive Emergency Checklist

## HYPERTENSIVE EMERGENCY:

- Two severe BP values ( $\geq 160/110$ ) taken 15-60 minutes apart. Values do not need to be consecutive.
- May treat within 15 minutes if clinically indicated

- Call for Assistance
- Designate:
  - Team leader
  - Checklist reader/recorder
  - Primary RN
- Ensure side rails up
- Administer seizure prophylaxis (magnesium sulfate first line agent, unless contraindicated)
- Antihypertensive therapy within 1 hour for persistent severe range BP
- Place IV; Draw preeclampsia labs
- Antenatal corticosteroids (if  $<34$  weeks of gestation)
- Re-address VTE prophylaxis requirement
- Place indwelling urinary catheter
- Brain imaging if unremitting headache or neurological symptoms
- Debrief patient, family, and obstetric team

## MAGNESIUM SULFATE

Contraindications: pulmonary edema, renal failure, myasthenia gravis

### IV access:

- Load 4-6 grams 10% magnesium sulfate in 100 mL solution over 20 min
- Label magnesium sulfate; Connect to labeled infusion pump
- Magnesium sulfate maintenance 1-2 grams/hour

### No IV access:

- 10 grams of 50% solution IM (5 g in each buttock)

## ANTIHYPERTENSIVE MEDICATIONS

For SBP  $\geq 160$  or DBP  $\geq 110$

- Labetalol** (20 mg, 40, 80 IV\* over 2 min, escalating doses, repeat q 10 min); Avoid in asthma or heart failure
- Hydralazine** (5-10 mg IV\* over 2 min, repeat q 20 min until target BP reached)
- Oral Nifedipine** (10, 20, 40 mg capsules; repeat BP q 20 min until target BP reached); Capsules should be administered orally, not punctured or otherwise administered sublingually

\* Maximum cumulative IV-administered doses should not exceed 220 mg labetalol or 25 mg hydralazine in 24 hours

**Note:** If first line agents unsuccessful, emergency consult with specialist (MFM, internal medicine, OB anesthesiology, critical care) is recommended

## ANTICONVULSANT MEDICATIONS

For recurrent seizures or when magnesium sulfate contraindicated

- Lorazepam (Ativan):** 2-4 mg IV x 1, may repeat once after 10-15 min
- Diazepam (Vallium):** 5-10 mg IV q 5-10 min to maximum dose 30 mg

EXAMPLE

13

- ✓ Call for assistance
- ✓ Designate team leader, checklist reader, primary RN
- ✓ Ensure side rails are up
- ✓ Administer seizure prophylaxis
- ✓ Antihypertensive therapy within 1 hr for persistent severe range BP
- ✓ Place IV; Draw PEC labs
- ✓ Antenatal corticosteroids is  $<34$  wks gestation
- ✓ Re-address VTE prophylaxis requirement
- ✓ Place indwelling urinary catheter
- ✓ Brain imaging if unremitting headache or neurological symptoms
- ✓ Debrief patient, family, OB team

# Eclampsia Checklist

EXAMPLE

14

- Call for Assistance
- Designate
  - Team leader
  - Checklist reader/recorder
  - Primary RN
- Ensure side rails up
- Protect airway and improve oxygenation:
  - Maternal pulse oximetry
  - Supplemental oxygen (100% non-rebreather)
    - Lateral decubitus position
    - Bag-mask ventilation available
    - Suction available
- Continuous fetal monitoring
- Place IV; Draw preeclampsia labs
- Administer magnesium sulfate
- Administer antihypertensive therapy if appropriate
- Develop delivery plan, if appropriate
- Debrief patient, family, and obstetric team

## MAGNESIUM SULFATE

Contraindications: pulmonary edema, renal failure, myasthenia gravis

### IV access:

- Load 4-6 grams 10% magnesium sulfate in 100 mL solution over 20 min
- Label magnesium sulfate; Connect to labeled infusion pump
- Magnesium sulfate maintenance 1-2 grams/hour

### No IV access:

- 10 grams of 50% solution IM (5 g in each buttock)

## ANTIHYPERTENSIVE MEDICATIONS

For SBP  $\geq$  160 or DBP  $\geq$  110

- Labetalol** (20 mg, 40, 80 IV\* over 2 min, escalating doses, repeat q 10 min); Avoid in asthma or heart failure, can cause neonatal bradycardia
- Hydralazine** (5-10 mg IV\* over 2 min, repeat q 20 min until target BP reached)

\* Maximum cumulative IV-administered doses should not exceed 220 mg labetalol or 25 mg hydralazine in 24 hours

**Note:** If persistent seizures, consider anticonvulsant medications and additional workup

## ANTICONVULSANT MEDICATIONS

For recurrent seizures or when magnesium sulfate contraindicated

- Lorazepam (Ativan):** 2-4 mg IV x 1, may repeat once after 10-15 min
- Diazepam (Valium):** 5-10 mg IV q 5-10 min to maximum dose 30 mg

## FOR PERSISTENT SEIZURES

- Neuromuscular block and intubate
- Obtain radiographic imaging
- ICU admission
- Consider anticonvulsant medications

- ✓ Call for assistance
- ✓ Designate team leader, checklist reader, primary RN
- ✓ Ensure side rails are up
- ✓ Protect airway + improve oxygenation
- ✓ Continuous fetal monitoring
- ✓ Place IV; Draw PEC labs
- ✓ Administer antihypertensive therapy if appropriate
- ✓ Develop delivery plan
- ✓ Debrief patient, family, OB team

# COMPLICATIONS & ESCALATION PROCESS

## MATERNAL (pregnant or postpartum)

- CNS (seizure, unremitting headache, visual disturbance)
- Pulmonary edema or cyanosis
- Epigastric or right upper quadrant pain
- Impaired liver function
- Thrombocytopenia
- Hemolysis
- Coagulopathy
- Oliguria \**<30 ml/hr for 2 consecutive hours*

## FETAL

- Abnormal fetal tracing
- IUGR

→ **Prompt evaluation and communication:** If undelivered, plan for delivery

# MONITORING CHANGE OF STATUS

*Once patient is stabilized, consider:*

## SEIZURE PROPHYLAXIS

- Magnesium sulfate (if not already initiated)

## TIMING & ROUTE OF DELIVERY

- **Eclampsia** → Delivery after stabilization
- **HELLP/Severe preeclampsia/Chronic hypertension + superimposed preeclampsia** → Vaginal delivery, if attainable in reasonable amount of time
- **≥ 34 weeks** → Deliver

## MATERNAL BP

- Continue control with oral agents
- Target range of 140-150/90-100

## IF PRETERM (<34 WKS) & EXPECTANT MGMT PLANNED

- Antenatal corticosteroids
- Subsequent pharmacotherapy
- **HELLP (Gestational age of fetal viability to 33 6/7 wks)**
- ✓ Delay delivery for 24-48 hours if maternal and fetal condition remains stable
- ✓ Contraindications to delay in delivery for fetal benefit of corticosteroids:
  - Uncontrolled hypertension
  - Eclampsia
  - Pulmonary edema
  - Suspected abruption placenta
  - Disseminated intravascular coagulation,
  - Nonreassuring fetal status
  - Intrauterine fetal demise

# GUIDELINES FOR DOCUMENTATION

EXAMPLE

17

ON ADMISSION	ASSESSMENT & PLAN
<ul style="list-style-type: none"><li>✓ <b>Complete history</b></li><li>✓ <b>Complete physical exam + preeclampsia symptoms:</b><ul style="list-style-type: none"><li>○ Unremitting headaches</li><li>○ Visual changes</li><li>○ Epigastric pain</li><li>○ Fetal activity</li><li>○ Vaginal bleeding</li></ul></li><li>✓ <b>Baseline BPs throughout pregnancy</b></li><li>✓ <b>Meds/drugs throughout pregnancy (illicit &amp; OTC)</b></li><li>✓ <b>Current vital signs, inc. O2 saturation</b></li><li>✓ <b>Current and past fetal assessment:</b><ul style="list-style-type: none"><li>○ FHR monitoring results</li><li>○ Est. fetal weight</li><li>○ BPP, as appropriate</li></ul></li></ul>	<ul style="list-style-type: none"><li>✓ <b>Indicate diagnosis of preeclampsia</b><ul style="list-style-type: none"><li>○ If no dx, indicate steps taken to exclude preeclampsia</li></ul></li><li>✓ <b>Antihypertensives taken (if any)</b><ul style="list-style-type: none"><li>○ Specific medications</li><li>○ Dose, route, frequency</li><li>○ Current fetal status</li></ul></li><li>✓ <b>Magnesium sulfate (if initiated for seizure prophylaxis)</b><ul style="list-style-type: none"><li>○ Dose, route, duration of therapy</li></ul></li><li>✓ <b>Delivery assessment</b><ul style="list-style-type: none"><li>○ If indicated, note: timing, method, route</li><li>○ If not indicated, describe circumstances to warrant delivery</li></ul></li><li>✓ <b>Antenatal corticosteroids if &lt; 34 weeks of gestation</b></li></ul> <div data-bbox="987 1258 1866 1368" style="border: 1px solid blue; padding: 5px;"><p><b>NOTE:</b> Continue ongoing documentation every 30 min until patient stabilized at &lt; SBP 160 or DBP 110</p></div>

# POSTPARTUM SURVEILLANCE

EXAMPLE

18

Necessary to prevent additional morbidity as preeclampsia/eclampsia can develop postpartum

## INPATIENT

- Measure BP every 4 hours after delivery until stable
- Do not use NSAIDs for women with elevated BP
- Do not discharge patient until BP is well controlled for at least 24 hours

## OUTPATIENT

- For pts with preeclampsia, visiting nurse evaluation recommended:
  - ✓ Within 3-5 days
  - ✓ Again in 7-10 days after delivery (earlier if persistent symptoms)

## ANTIHYPERTENSIVE THERAPY

- Recommended for persistent postpartum HTN: SBP  $\geq$  150 or DBP  $\geq$  100 on at least two occasions at least 4 hours apart
- Persistent SBP  $\geq$  160 or DBP  $\geq$  110 should be treated within 1 hour

# Postpartum Preeclampsia Checklist

## IF PATIENT < 6 WEEKS POSTPARTUM WITH:

- BP  $\geq$  160/110 or
- BP  $\geq$  140/90 with unremitting headache, visual disturbances, epigastric pain

- Call for Assistance
- Designate:
  - Team leader
  - Checklist reader/recorder
  - Primary RN
- Ensure side rails up
- Call obstetric consult; Document call
- Place IV; Draw preeclampsia labs
  - CBC
  - Chemistry Panel
  - PT
  - Uric Acid
  - PTT
  - Hepatic Function
  - Fibrinogen
  - Type and Screen
- Administer seizure prophylaxis
- Administer antihypertensive therapy
  - Contact MFM or Critical Care for refractory blood pressure
- Consider indwelling urinary catheter
  - Maintain strict I&O - patient at risk for pulmonary edema
- Brain imaging if unremitting headache or neurological symptoms

### MAGNESIUM SULFATE

Contraindications: pulmonary edema, renal failure, myasthenia gravis

#### IV access:

- Load 4-6 grams 10% magnesium sulfate in 100 mL solution over 20 min
- Label magnesium sulfate; Connect to labeled infusion pump
- Magnesium sulfate maintenance 1-2 grams/hour

#### No IV access:

- 10 grams of 50% solution IM (5 g in each buttock)

### ANTIHYPERTENSIVE MEDICATIONS

For SBP  $\geq$  160 or DBP  $\geq$  110

- Labetalol** (20 mg, 40, 80 IV\* over 2 min, escalating doses, repeat q 10 min); Avoid in asthma or heart failure
- Hydralazine** (5-10 mg IV\* over 2 min, repeat q 20 min until target BP reached)
- Oral Nifedipine** (10, 20, 40 mg capsules; repeat BP q 20 min until target BP reached); Capsules should be administered orally, not punctured or otherwise administered sublingually

\* Maximum cumulative IV-administered doses should not exceed 220 mg labetalol or 25 mg hydralazine in 24 hours

**Note:** If first line agents unsuccessful, emergency consult with specialist (MFM, internal medicine, OB anesthesiology, critical care) is recommended

### ANTICONVULSANT MEDICATIONS

For recurrent seizures or when magnesium sulfate contraindicated

- Lorazepam (Ativan):** 2-4 mg IV x 1, may repeat once after 10-15 min
- Diazepam (Valium):** 5-10 mg IV q 5-10 min

- ✓ Call for assistance
- ✓ Designate team leader, checklist reader, primary RN
- ✓ Ensure side rails up
- ✓ Call OB consult; Document call
- ✓ Place IV; Draw PEC labs
- ✓ Administer seizure prophylaxis
- ✓ Administer antihypertensive therapy
- ✓ Consider indwelling urinary catheter. Maintain strict I&O
- ✓ Brain imaging if unremitting headache or neurological symptoms

# DISCHARGE PLANNING

## All patients receive information on preeclampsia:

- ✓ Signs and symptoms
- ✓ Importance of reporting information to health care provider as soon as possible
- ✓ Culturally-competent, patient-friendly language

## All new nursing and physician staff receive information on hypertension in pregnancy and postpartum

### FOR PATIENTS WITH PREECLAMPSIA

---

- ✓ BP monitoring recommended 72 hours after delivery
- ✓ Outpatient surveillance (visiting nurse evaluation) recommended:
  - Within 3-5 days
  - Again in 7-10 days after delivery (earlier if persistent symptoms)

# POST-DISCHARGE EVALUATION

EXAMPLE

21

## ELEVATED BP AT HOME, OFFICE, TRIAGE

### Postpartum triggers:

- SBP  $\geq$  160 or DBP  $\geq$  110 or
- SBP  $\geq$  140-159 or DBP  $\geq$  90-109 with unremitting headaches, visual disturbances, or epigastric/RUQ pain



- Emergency Department treatment (OB /MICU consult as needed)
- AntiHTN therapy suggested if persistent **SBP  $\geq$  150 or DBP  $\geq$  100** on at least two occasions at least 4 hours apart
- Persistent **SBP  $\geq$  160 or DBP  $\geq$  110** should be treated within 1 hour



Good response to antiHTN treatment and asymptomatic



Signs and symptoms of eclampsia, abnormal neurological evaluation, congestive heart failure, renal failure, coagulopathy, poor response to antihypertensive treatment



Admit for further observation and management (L&D, ICU, unit with telemetry)



Recommend emergency consultation for further evaluation (MFM, internal medicine, OB anesthesiology, critical care)



# CONCLUSION

- **Systolic BP  $\geq$  160 or diastolic BP  $\geq$  110** warrant:
  - ✓ Prompt evaluation at bedside
  - ✓ Treatment to decrease maternal morbidity and mortality
  
- Risk reduction and successful clinical outcomes require avoidance/management of severe systolic and diastolic hypertension in women with:
  - ✓ Preeclampsia
  - ✓ Eclampsia
  - ✓ Chronic hypertension + superimposed preeclampsia
  
- Increasing evidence indicates that standardization of care improves patient outcomes

# Index of Hypertension Bundle Content Changes (November 2015)

## EDUCATIONAL MATERIALS

### Severe Hypertension Slide Deck

Slide 5	<b>REVISED</b> – Definition of hypertensive emergency
6	<b>REVISED</b> – When to treat following: <ul style="list-style-type: none"> <li>• One severe hypertensive value</li> <li>• Hypertensive emergency</li> </ul>
7	<b>ADDED</b> <ul style="list-style-type: none"> <li>• Oral nifedipine as first line therapy option</li> <li>• Mag sulfate and anticonvulsant recommendations</li> </ul>
8	<b>ADDED</b> <ul style="list-style-type: none"> <li>• BP check 10 min after second 80 mg dose of labetalol</li> <li>• Language on adverse effects and use of clinical judgement</li> </ul>
9	<b>REVISED</b> - BP check timeframes <b>ADDED</b> - Language on adverse effects and use of clinical judgement
10	<b>ADDED</b> – Oral nifedipine algorithm
11	<b>ADDED</b> – <ul style="list-style-type: none"> <li>• Use of oral nifedipine algorithm if no IV access</li> <li>• Second line therapies: Labetalol or nicardipine infusion pump, sodium nitroprusside</li> </ul>
13 – 14, 19	<b>REVISED</b> – Streamlined format
16	<b>ADDED</b> – Target range for maternal BP
18	<b>ADDED</b> – Included under postpartum surveillance: <ul style="list-style-type: none"> <li>• Antihypertensive meds</li> <li>• Recommendations for outpatients w/ preeclampsia</li> </ul>

### Algorithms

(Labetalol, Hydralazine, Oral Nifedipine)

*Available online in printable, PDF format*

### Checklists

(Hypertensive Emergency, Eclampsia, Postpartum Preeclampsia – ED)

*Available online in printable, PDF format*

# References

- Abalos E, Duley L, Steyn DW, Henderson-Smart DJ. "Antihypertensive drug therapy for mild to moderate hypertension during pregnancy (review)." *The Cochrane Collaboration*. 2007, Issue 1. Art. No.: CD002252. DOI: 10.1002/14651858.CD002252.pub2.
- The American College of Obstetricians and Gynecologists. "Hypertension in Pregnancy." Task Force on Hypertension in Pregnancy, 2013.
- The American College of Obstetricians and Gynecologists. "Emergent Therapy for Acute-Onset, Severe Hypertension with Preeclampsia or Eclampsia." Committee Opinion, Number 514. 2011. <http://tinyurl.com/m93r3oc>
- Churchill D, Duley L. "Interventionist versus expectant care for severe pre-eclampsia before term." *The Cochrane Collaboration*. 2002, Issue 3. Art. No.: CD003106. DOI: 10.1002/14651858.CD003106.
- Duley L, Gülmezoglu AM, Henderson-Smart DJ. "Magnesium sulphate and other anticonvulsants for women with preeclampsia (review)." *The Cochrane Collaboration*. 2003, Issue 2. Art. No.: CD000025. DOI: 10.1002/14651858.CD000025.
- Duley L, Henderson-Smart DJ, Meher S. "Drugs for treatment of very high blood pressure during pregnancy (review)." *The Cochrane Collaboration*. 2006, Issue 3. Art. No.: CD001449. DOI: 10.1002/14651858. CD001449.pub2.
- Maurice L. Druzin, MD; Laurence E. Shields, MD; Nancy L. Peterson, RNC, PNNP, MSN; Valerie Cape, BSBA. "Preeclampsia Toolkit: Improving Health Care Response to Preeclampsia." California Maternal Quality Care Collaborative Toolkit to Transform Maternity Care. Developed under contract #11-10006 with the California Department of Public Health; Maternal, Child and Adolescent Health Division; Published by the California Maternal Quality Care Collaborative, November 2013.
- National Institute for Health and Clinical Excellence. "The management of hypertensive disorders during pregnancy." *NICE Clinical Guideline #107*. Modified January 2011.
- New York State Department of Health. "Hypertensive Disorders in Pregnancy." *NYSDOH Executive - Guideline Summary*, May 2013.
- Shekhar et al. "Oral Nifedipine or Intravenous Labetalol for Hypertensive Emergency in Pregnancy." *Obstetrics and Gynecology*, 2012 (122): 1057-1063.
- Sibai BM. "Etiology and management of postpartum hypertension-preeclampsia." *American Journal of Obstetrics and Gynecology*. 2012: 470-5.
- Smith M, Waugh J, Nelson-Piercy C. "Management of postpartum hypertension." *The Obstetrician & Gynaecologist*, 2013: 15:45-50.
- WHO Recommendations for the prevention and treatment of pre-eclampsia and eclampsia. *World Health Organization, 2011*. Geneva, Switzerland.